

© Engineered Science Publisher LLC 2021

# **Supporting Information**

## **Maxwell-Wagner Relaxations in Ca-, Sm- and Nd-doped Ceria**

R. Ahmed<sup>1</sup>, S. Renjun<sup>1</sup>, S. ur Rehman<sup>2</sup>, T. Y. Li<sup>1</sup>, Y. Yu<sup>1</sup>, Q. J. Li<sup>1</sup>, C. C. Wang<sup>1,3\*</sup>

<sup>1</sup> Laboratory of Dielectric Functional Materials, School of Physics & Material Science,  
Anhui University, Hefei, 230601, China

<sup>2</sup> High Magnetic Field Laboratory, Chinese Academy of Sciences, Hefei 230031, Anhui, P.  
R. China.

<sup>3</sup> State Key Laboratory of Low-Dimensional Quantum Physics, Department of Physics, Tsinghua  
University, Beijing 100084, China

Email: ccwang@ahu.edu.cn




Figure S1. Two-relaxation fittings for the electric modulus spectra of CSO (a-g) and the resultant Arrhenius plots of the relaxations (h).




Figure S2. Two-relaxation fittings for the electric modulus spectra of CNO (a-g) and the resultant Arrhenius plots of the relaxations (h).