

Volatile Components in the Extracts from 20 Herbaceous Plant Flowers

Limin Wang^{1,#}, Wan Adibah Wan Mahari^{2,#}, Rock Keey Liew^{3,#}, Guanyan Li¹, Xiaochen Yue^{1,*},
Haiping Gu¹, Wanxi Peng^{1,4,*} and Xiangmeng Chen^{1,*}

¹College of Forestry, College of Science, College of Landscape Architecture and Art, Henan Agricultural University, Zhengzhou 450002, China

²Higher Institution Centre of Excellence (HICoE), Institute of Tropical Aquaculture and Fisheries, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia

³Center for Global Health Research (CGHR), Saveetha Medical College, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Chennai, India

⁴Longzihu New Energy Laboratory, Zhengzhou Institute of Emerging Industrial Technology, Zhengzhou, 450000, China

[#]These authors contributed equally to this work.

*Email: yuxiaochen95@163.com (Xiaochen Yue); wanxipeng2024@163.com (Wanxi Peng); xmchen0610@163.com (Xiangmeng Chen)

DOI:<https://dx.doi.org/10.30919/faf1612>

Proof of authorization for Tables




Fig S1: Total ion chromatograms of GC-MS for the ethanol extracts of the flowers of *Taraxacum mongolicum* Hand.-Mazz.

Table S1: GC-MS analysis result of ethanol extract from *Taraxacum mongolicum* Hand.-Mazz.

Number	materials	relative content
1	Dicyclopentadiene diepoxide	0.23%
2	Methylephedrine	4.18%
3	3-trsns-(1,1-dimethylethyl)-4-trans-methoxycyclohexanol	1.49%
4	Benzoic acid, 2-(dimethylamino)ethyl ester	1.38%
5	6-Hepten-3-one, 5-hydroxy-4-methyl-	2.84%
6	(Methoxymethyl)trimethylsilane	3.67%
7	(2-Methyl-1-cyclohexenyl)methanol, TMS derivative	0.25%
8	4-Cyclohexene-1,2-diol	0.29%
9	Decanoic acid, 3-hydroxy-, methyl ester	0.52%
10	Dodecane, 1-fluoro-	0.88%
11	5-Hepten-2-amine, N,6-dimethyl-	3.84%
12	1-Deoxy-d-mannitol	6.48%
13	5,7-Dodecadiyn-1,12-diol	0.19%
14	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	0.32%
15	3-n-Butylthiolane	1.54%
16	Melezitose	2.21%
17	4-Cyanothiophenol	3.43%
18	7-Ethyl-4-decen-6-one	6.63%
19	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	0.37%
20	2-Cyclohexylpiperidine	0.82%
21	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	6.12%
22	Ethyl stearate, 9,12-diepoxo	0.99%
23	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	1.68%
24	Triisobutyl(3-phenylpropoxy)silane	0.65%
25	Tetradecanoic acid	5.83%
26	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	0.18%
27	Hexadecanoic acid, methyl ester	2.96%
28	n-Hexadecanoic acid	29.92%
29	Estra-1,3,5(10)-trien-17.beta.-ol	10.09%


**Fig S2:** Total ion chromatograms of GC-MS for the ethanol extracts of the flowers of *Cirsium arvense* var. *integrifolium* Wimm. & Grab.

Table S2: GC-MS analysis result of ethanol extract of *Cirsium arvense* var. *integrifolium* Wimm. & Grab.

Number	materials	relative content
1	Cyclotrisiloxane, hexamethyl-	0.31%
2	Trifluoroethanol, TBDMS derivative	3.55%
3	Benzamide, N-[6-(2-furyl)-2-oxo-2H-pyran-3-yl]-	1.96%
4	Benzoyl isothiocyanate	2.29%
5	1,3-Propanediol, 2-methyl-, dipropanoate	1.16%
6	Methyl 12,13-tetradecadienoate	0.83%
7	Dodecane, 1-fluoro-	4.64%
8	15-Tetracosenoic acid, methyl ester, (Z)-	43.87%
9	9,12-Octadecadienoic acid (Z,Z)-	13.80%
10	9,12-Octadecadienoyl chloride, (Z,Z)-	1.53%
11	trans-13-Octadecenoic acid	2.36%
12	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	0.68%
13	Tetracosanoic acid, methyl ester	6.39%
14	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	11.01%
15	tert-Hexadecanethiol	1.40%
16	Estra-1,3,5(10)-trien-17.beta.-ol	3.86%
17	Glycidyl (Z)-9-Heptadecenoate	0.36%




Fig S3: Total ion chromatograms of GC-MS for the ethanol extracts of the flowers of *Cyanus segetum* Hill.

Table S3: GC-MS analysis result of ethanol extract of *Cyanus segetum* Hill.

Number	materials	relative content
1	Trifluoroethanol, TBDMS derivative	11.99%
2	Methanone, (4-bromo-5-methyl-2-nitro-3-thienyl)(phenyl)-	0.71%
3	1,3-Propanediol, TMS derivative	0.34%
4	Benzoic acid, 2-(dimethylamino)ethyl ester	1.36%
5	Methcathinone	1.75%
6	Ethanol, TMS derivative	6.31%
7	1,5-Anhydro-d-mannitol	0.50%
8	Dodecane, 1-fluoro-	5.46%
9	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	5.67%
10	3-n-Butylthiolane	3.50%
11	Melezitose	1.57%
12	Methyl 4,7,10,13-hexadecatetraenoate	1.46%
13	2,5-Octadecadiynoic acid, methyl ester	0.90%
14	tert-Hexadecanethiol	2.98%
15	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	20.16%
16	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	1.77%
17	5,8,11-Eicosatrienoic acid, (Z)-, TMS derivative	1.22%
18	Estra-1,3,5(10)-trien-17.beta.-ol	14.53%
19	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	16.64%
20	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	1.18%




Fig S4: Total ion chromatograms of GC-MS for the ethanol of *Hemisteptia lyrata* (Bunge) Fisch. & C. A. Mey.

Table S4: GC-MS analysis result of ethanol extract of *Hemisteptia lyrata* (Bunge) Fisch. & C. A. Mey.

Number	materials	relative content
1	Dicyclopentadiene diepoxide	2.16%
2	Perfluoropropionic acid, TBDMS derivative	7.20%
3	Benzoic acid, 2-(dimethylamino)ethyl ester	0.54%
4	Ephedrine	0.55%
5	1,3-Propanediol, TMS derivative	0.64%
6	3-(Methylthio)propyl benzoate	4.39%
7	1,3-Propanediol, 2-methyl-, dipropionate	3.20%
8	Propanoic acid, butyl ester	28.28%
9	.beta.-D-Glucopyranoside, methyl	1.40%
10	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	2.03%
11	3-n-Butylthiolane	2.18%
12	2,2-Dimethyl-5-[2-(2-trimethylsilylethoxymethoxy)-propyl]-[1,3]dioxolane-4-carboxaldehyde	0.20%
13	<i>Caryophyllene</i>	6.43%
14	2,5-Octadecadiynoic acid, methyl ester	2.72%
15	tert-Hexadecanethiol	1.59%
16	5,8,11-Eicosatrienoic acid, (Z)-, TMS derivative	0.21%
17	.alpha.-Linolenic acid, TMS derivative	1.29%
18	2,5-Octadecadiynoic acid, methyl ester	1.13%

Number	materials	relative content
19	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	11.55%
20	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	1.36%
21	Estra-1,3,5(10)-trien-17.beta.-ol	13.80%
22	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	5.03%
23	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	2.12%




Fig S5: Total ion chromatograms of GC-MS for the ethanol of *Coreopsis basalis* (A. Dietr.) S. F. Blake.

Table S5: GC-MS analysis result of ethanol extract of *Coreopsis basalis* (A. Dietr.) S. F. Blake

Number	materials	relative content
1	Perfluoropropionic acid, TBDMS derivative	2.26%
2	1,3-Propanediol, TMS derivative	1.37%
3	Undec-10-ynoic acid, heptyl ester	0.30%
4	3-n-Butylthiolane	0.84%
5	1,3-Disilacyclobutane, 1,1,3,3-tetramethyl-	0.38%
6	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	0.89%
7	7-Ethyl-4-decen-6-one	0.74%
8	Methyl 2,4-tridecadiynoate	0.16%
9	2,5-Octadecadiynoic acid, methyl ester	3.82%
10	Melezitose	0.25%

Number	materials	relative content
11	2,2-Dimethyl-5-[2-(2-trimethylsilylethoxymethoxy)-propyl]-[1,3]dioxolane-4-carboxaldehyde	0.22%
12	((1R,2S,6S,7S,8S)-8-Isopropyl-1-methyl-3-methylenetricyclo[4.4.0.0 ^{2,7}]decane-rel-	31.29%
13	.beta.-Longipinene	0.75%
14	cis-.alpha.-Bergamotene	0.92%
15	Methyl 4,7,10,13-hexadecatetraenoate	0.33%
16	Methyl 3,5-tetradecadiynoate	0.29%
17	Ylangene	0.40%
18	<i>Benzene, 1-(1,5-dimethyl-4-hexenyl)-4-methyl-</i>	3.44%
19	<i>.beta.-Bisabolene</i>	9.74%
20	Bicyclo[4.4.0]dec-1-ene, 2-isopropyl-5-methyl-9-methylene-	0.21%
21	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	1.07%
22	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	5.62%
23	5,8,11-Eicosatrienoic acid, (Z)-, TMS derivative	0.36%
24	Benzene, 1,3,5-heptatriyn-1-yl-	18.77%
25	3-(1,5-Dimethyl-hex-4-enyl)-2,2-dimethyl-cyclopent-3-enol	7.13%
26	Dibenzothiophene, 4-methyl-	4.55%
27	1,25-Dihydroxyvitamin D3, TMS derivative三烯-1,3β-二醇	0.66%
28	Estra-1,3,5(10)-trien-17.beta.-ol	1.77%
29	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	1.11%
30	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	0.35%




Fig S6: Total ion chromatograms of GC-MS for the ethanol of *Trifolium repens* L.

Table S6: GC-MS analysis result of ethanol extract from *Trifolium repens* L.

Number	materials	relative content
1	Methanol, tris(methylenecyclopropyl)-	5.93%
2	Methanone, (4-bromo-5-methyl-2-nitro-3-thienyl)(phenyl)-	0.32%
3	S(-)-Methcathinone	0.36%
4	Benzoyl isothiocyanate	1.70%
5	.beta.-D-Glucopyranose, 1-thio-1-[N-hydroxy-5-(methylthio)pentanimidate]	0.26%
6	2,5-Octadecadiynoic acid, methyl ester	0.22%
7	.alpha.-Linolenic acid, TMS derivative	1.37%
8	Linoleic acid ethyl ester	2.33%
9	2,5-Octadecadiynoic acid, methyl ester	1.12%
10	Triisobutyl(3-phenylpropoxy)silane	0.79%
11	3,7,11,14,18-Pentaoxa-2,19-disilaecosane, 2,2,19,19-tetramethyl-	72.44%
12	3,7,11,14,18-Pentaoxa-2,19-disilaecosane, 2,2,19,19-tetramethyl-	0.53%
13	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	3.18%
14	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	5.95%
15	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	3.49%




Fig S7: Total ion chromatograms of GC-MS for the ethanol of *Paeonia lactiflora* Pall.

Table S7: GC-MS compound analysis of ethanol extract from *Paeonia lactiflora* Pall.

Number	materials	relative content
1	Perfluoropropionic acid, TBDMS derivative	9.02%
2	Benzamide, N-[6-(2-furyl)-2-oxo-2H-pyran-3-yl]-	2.32%
3	1-Propanone, 2-bromo-1-phenyl-	0.08%
4	1,3-Propanediol, TMS derivative	0.84%
5	3,5-Dimethylpyrazole	1.18%
6	trans-4-Methylcyclohexanol, trifluoroacetate	0.59%
7	4-Methylcyclohexaneacetic acid (stereoisomer 1)	0.58%
8	2,5,6-Trimethyl-1,3-oxathiane	0.37%
9	3-n-Butylthiolane	19.25%
10	Phenyl-.beta.-D-glucoside	0.30%
11	2,5,6-Trimethyl-1,3-oxathiane	0.91%
12	Lactose	1.31%
13	Ethylene glycol, O,O-di(pivaloyl)-	0.63%
14	Pyrovalerone	5.05%
15	Methyl 6-oxoheptanoate	1.40%
16	Hexanediamide, N,N'-di-benzoyloxy-	6.62%
17	.beta.-D-Glucopyranose, 1-thio-, 1-[N-hydroxy-5-(methylthio)pentanimidate]	1.95%
18	4-Mercaptophenol	5.91%

Number	materials	relative content
19	Melezitose	4.36%
20	Methyl 3,5-tetradecadiynoate	0.09%
21	Gallacetophenone-4'-methylether	0.10%
22	Resorcinol, TMS derivative	11.93%
23	Gallacetophenone-4'-methylether	7.60%
24	4-Hexenoic acid, 4-methyl-6-(fluorodimethylsilyl)-6-trimethylsilyl-	0.37%
25	5,8,11-Eicosatriynoic acid, tert-butyldimethylsilyl ester	0.15%
26	.alpha.-D-Galactopyranose, 6-O-(trimethylsilyl)-, cyclic 1,2:3,4-bis(methylboronate)	0.57%
27	.alpha.-D-Galactopyranose, 6-O-(trimethylsilyl)-, cyclic 1,2:3,4-bis(methylboronate)	7.25%
28	5,8,11,14-Eicosatetraynoic acid, TMS derivative	0.40%
29	(1,25-Dihydroxyvitamin D3, TMS derivative	1.35%
30	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	7.51%




Fig S8: Total ion chromatograms of GC-MS for the ethanol of *Erigeron annuus* (L.) Pers.

Table S8: GC-MS analysis result of ethanol extract of *Erigeron annuus* (L.) Pers.

Number	materials	relative content
1	Perfluoropropionic acid, TBDMS derivative	12.14%
2	Methanone, (4-bromo-5-methyl-2-nitro-3-thienyl)(phenyl)-	1.28%
3	Benzoyl isothiocyanate	0.49%
4	Ephedrine	0.53%

Number	materials	relative content
5	Ethanol, TMS derivative	7.13%
6	2-Dimethylsilyloxytridecane	0.10%
7	Lactose	3.35%
8	Arachidonic acid, TMS derivative	0.59%
9	10-Undecynoic acid, TMS derivative	0.22%
10	Methyl 4,7,10,13-hexadecatetraenoate	1.36%
11	2,5-Octadecadiynoic acid, methyl ester	5.10%
12	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	4.52%
13	(1R,2S,6S,7S,8S)-8-Isopropyl-1-methyl-3-methylenetricyclo[4.4.0.0 ^{2,7}]decane-rel-	1.72%
14	Methyl 4,7,10,13-hexadecatetraenoate	2.61%
15	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	20.09%
16	1,25-Dihydroxyvitamin D3, TMS derivative	2.92%
17	10-Undecynoic acid, TMS derivative	13.05%
18	1-Heptatriacotanol	1.35%
19	tert-Hexadecanethiol	8.42%
20	1,25-Dihydroxyvitamin D3, TMS derivative	2.13%
21	Estra-1,3,5(10)-trien-17.β.-ol	5.63%
22	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	3.65%
23	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	1.62%




Fig S9: Total ion chromatograms of GC-MS for the ethanol of *Oxalis corymbosa* DC.

Table S9: GC-MS analysis result of ethanol extract from *Oxalis corymbosa* DC.

Number	materials	relative content
1	Trifluoromethyltrimethylsilane	12.74%
2	Perfluoropropionic acid, TBDMS derivative	1.55%
3	3-(6,7-Dimethoxy-2H-1,3-benzodioxol-5-yl)-1-phenylprop-2-en-1-one	3.05%
4	Pseudoephedrine	0.77%
5	Benzoyl isothiocyanate	2.09%
6	2-Amino-2-methyl-1,3-propanediol	17.14%
7	3-n-Butylthiolane	1.67%
8	2,2-Dimethyl-5-[2-(2-trimethylsilylethoxymethoxy)-propyl]-[1,3]dioxolane-4-carboxaldehyde	0.44%
9	2,2-Dimethyl-5-[2-(2-trimethylsilylethoxymethoxy)-propyl]-[1,3]dioxolane-4-carboxaldehyde	5.79%
10	Melezitose	1.76%
11	Ethyl pipecolate	7.49%
12	2-Cyclohexylpiperidine	1.45%
13	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	6.47%
14	1,25-Dihydroxyvitamin D3, TMS derivative	0.99%
15	tert-Hexadecanethiol	6.18%
16	Arachidonic Acid, TBDMS derivative	0.63%
17	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	8.18%
18	Hexadecanoic acid, ethyl ester	21.34%
19	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	0.27%




Fig S10: Total ion chromatograms of GC-MS for the ethanol of *Hemerocallis fulva* 'Golden Doll '

Table S10: GC-MS analysis result of ethanol extract from *Hemerocallis fulva* 'Golden Doll. '

Number	materials	relative content
1	Trifluoroethanol, TBDMS derivative	10.14%
2	3-Hexen-1-ol benzoate	3.16%
3	Benzoic acid, 2-(dimethylamino)ethyl ester	1.66%
4	6-Hepten-3-one, 5-hydroxy-4-methyl-	0.42%
5	1,3-Propanediol, TMS derivative	0.35%
6	1H-Pyrazole, 1,5-dimethyl-	0.27%
7	3-Furanmethanol	6.17%
8	3-n-Butylthiolane	7.12%
9	Dodecane, 1-fluoro-	0.56%
10	Cholestan-22(26)-epoxy-3,16-dione	0.69%
11	Methyl 6-oxoheptanoate	0.09%
12	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	4.70%
13	7-Ethyl-4-decen-6-one	6.69%
14	t-Butyl(4,5-dihydrofuran-2-yl)dimethylsilane	0.60%
15	2,2-Dimethyl-5-[2-(2-trimethylsilylethoxymethoxy)-propyl]-[1,3]dioxolane-4-carboxaldehyde	0.34%
16	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	0.15%
17	Melezitose	6.63%

Number	materials	relative content
18	Ethyl 9.cis.,11.trans.-octadecadienoate	15.05%
19	trans-13-Octadecenoic acid	3.39%
20	.alpha.-Linolenic acid, TMS derivative	6.86%
21	9,12-Octadecadiynoic acid, tert-butyldimethylsilyl ester	1.36%
22	Octadecanoic acid, ethyl ester	3.19%
23	10-Undecynoic acid, TMS derivative	2.91%
24	1,25-Dihydroxyvitamin D3, TMS derivative	3.17%
25	Estra-1,3,5(10)-trien-17.beta.-ol	3.78%
26	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	9.31%
27	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	0.95%
28	Doconexent, TBDMS derivative	0.28%




Fig S11: Total ion chromatograms of GC-MS for the ethanol of *Hosta plantaginea* (Lam.) Aschers..

Table S11: GC-MS analysis result of ethanol extract of *Hosta plantaginea* (Lam.) Aschers..

Number	materials	relative content
1	Trifluoroethanol, TBDMS derivative	18.59%
2	3-Buten-2-ol, benzoate	0.60%
3	Benzenemethanol, .alpha.-[1-(methylamino)ethyl]-, [S-(R*,S*)]-	0.62%
4	Benzoyl isothiocyanate	5.16%
5	2,5,6-Trimethyl-1,3-oxathiane	6.26%
6	3-n-Butylthiolane	8.27%
7	<i>Alpha-l-rhamnopyranose</i>	0.70%
8	5-Thio-D-glucose	0.37%
9	Thymine	3.31%
10	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	17.01%
11	D-Fructose, diethyl mercaptal, pentaacetate	1.55%
12	2,2-Dimethyl-5-[2-(2-trimethylsilylethoxymethoxy)-propyl]-[1,3]dioxolane-4-carboxaldehyde	0.56%
13	Methyl 3,5-tetradecadiynoate	0.33%
14	Melezitose	9.89%
15	Triisobutyl(3-phenylpropoxy)silane	1.60%
16	10-Undecynoic acid, TMS derivative	3.96%
17	1,25-Dihydroxyvitamin D3, TMS derivative	1.99%
18	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	1.35%
19	Estra-1,3,5(10)-trien-17.beta.-ol	4.27%
20	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	12.99%
21	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	0.62%




Fig S12: Total ion chromatograms of GC-MS for the ethanol of *Setaria viridis* (L.) Beauv.

Table S12: GC-MS analysis result of ethanol extract from *Setaria viridis* (L.) Beauv.

Number	materials	relative content
1	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	10.44%
2	Benzamide, N-[6-(2-furyl)-2-oxo-2H-pyran-3-yl]-	0.62%
3	Benzoic acid, 2-(dimethylamino)ethyl ester	0.39%
4	Ephedrine	0.52%
5	1,3-Propanediol, TMS derivative	1.57%
6	1-Deoxy-d-mannitol	0.80%
7	Ethyl(dimethyl)ethoxysilane	3.44%
8	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	2.17%
9	2,5-Octadecadiynoic acid, methyl ester	2.76%
10	Thymidine	2.25%
11	Methyl 3,5-tetradecadiynoate	0.57%
12	Phenol, 3,5-bis(1,1-dimethylethyl)-	0.16%
13	Ethyl stearate, 9,12-diepoxy	0.59%
14	Undec-10-ynoic acid, butyl ester	0.73%
15	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	4.14%
16	1,25-Dihydroxyvitamin D3, TMS derivative	0.64%
17	tert-Hexadecanethiol	5.61%
18	n-Hexadecanoic acid	22.07%

Number	materials	relative content
19	Phthalic acid, butyl oct-3-yl ester	8.83%
20	Hexadecanoic acid, ethyl ester	31.70%




Fig S13: Total ion chromatograms of GC-MS for the ethanol of *Cyperus rotundus* L.

Table S13: GC-MS analysis result of ethanol extract from *Cyperus rotundus* L.

Number	materials	relative content
1	Hexadecanoic acid, ethyl ester	5.21%
2	Benzamide, N-[6-(2-furyl)-2-oxo-2H-pyran-3-yl]-	1.50%
3	1,3-Propanediol, TMS derivative	1.95%
4	1-Deoxy-d-mannitol	0.18%
5	Methyl 6-oxoheptanoate	0.72%
6	3-n-Butylthiolane	1.18%
7	2,5-Octadecadiynoic acid, methyl ester	1.56%
8	cis-5,8,11,14,17-Eicosapentaenoic acid	0.48%
9	Melezitose	2.04%
10	Methyl 4,7,10,13-hexadecatetraenoate	1.23%
11	1H-Cyclopropa[a]naphthalene, decahydro-1,1,3a-trimethyl-7-methylene-, [1aS-(1a.alpha.,3a.alpha.,7a.beta.,7b.alpha.)]-	3.42%
12	cis-13-Octadecenoic acid	10.70%
13	Ethyl 9.cis.,11.trans.-octadecadienoate	15.56%

Number	materials	relative content
14	(E)-9-Octadecenoic acid ethyl ester	30.61%
15	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	15.32%
16	Octadecanoic acid, ethyl ester	3.28%
17	Doconexent, TMS derivative	0.41%
18	Tricyclo[5.4.3.0(1,8)]tetradecan-6-one, 4-ethenyl-3-hydroxy-2,4,7,14-tetramethyl	4.65%




Fig S14: Total ion chromatograms of GC-MS for the ethanol of *Hemerocallis hybrida* Bergmans.

Table S14: GC-MS analysis result of ethanol extract from *Hemerocallis hybrida* Bergmans.

Number	materials	relative content
1	Perfluoropropionic acid, TBDMS derivative	14.69%
2	Furan, 3-methyl-	3.33%
3	Benzoic acid, 2-(dimethylamino)ethyl ester	3.34%
4	1,3-Propanediol, TMS derivative	1.58%
5	1,1-Cyclohexanedimethanol	0.45%
6	1H-Imidazole-4-methanol	14.69%
7	3-n-Butylthiolane	11.28%
8	Dodecane, 1-fluoro-	2.60%
9	Thymine	0.99%
10	tert-Nonyl mercaptan	0.63%

Number	materials	relative content
11	2-Hexene, 1,1-diethoxy-	0.51%
12	1-Methoxy-1-methyl-1-silacyclohexane	5.08%
13	4-Mercaptophenol	1.99%
14	Melezitose	4.98%
15	2-tert-Butyl-4-methylphenol, tert-butyldimethylsilyl ether	1.42%
16	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	3.07%
17	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	5.17%
18	1,25-Dihydroxyvitamin D3, TMS derivative	2.75%
19	Estra-1,3,5(10)-trien-17.β-ol	6.70%
20	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	14.46%
21	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	0.30%




Fig S15: Total ion chromatograms of GC-MS for the ethanol of *Echinacea purpurea*.

Table S15: GC-MS analysis result of ethanol extract of *Echinacea purpurea*.

Number	materials	relative content
1	Perfluoropropionic acid, TBDMS derivative	4.11%
2	Ephedrine	1.04%
3	1,3-Propanediol, TMS derivative	0.38%
4	.beta.-D-Glucopyranose, 1-thio-, 1-[N-hydroxy-5-(methylthio)pentanimidate]	1.43%
5	1-Deoxy-d-mannitol	1.61%
6	Copaene	1.16%
7	(1R,2S,6S,7S,8S)-8-Isopropyl-1-methyl-3-methylenetricyclo[4.4.0.02,7]decane-rel-	68.00%
8	cis-.alpha.-Bergamotene	1.56%
9	Tricyclo[4.4.0.0(2,7)]decane, 1-methyl-3-methylene-8-(1-methylethyl)-, stereoisomer	1.79%
10	<i>Caryophyllene</i>	2.52%
11	(1R,2S,6S,7S,8S)-8-Isopropyl-1-methyl-3-methylenetricyclo[4.4.0.02,7]decane-rel-	1.34%
12	Tau-Cadinol acetate	0.50%
13	Methyl 4,7,10,13-hexadecatetraenoate	0.44%
14	2,5-Octadecadiynoic acid, methyl ester	0.82%
15	(Cyclohexanol, 3-ethenyl-3-methyl-2-(1-methylethenyl)-6-(1-methylethyl)-, [1R-(1.alpha.,2.alpha.,3.beta.,6.alpha.)]-	0.65%
16	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	6.62%
17	(1R,7S,E)-7-Isopropyl-4,10-dimethylenecyclodec-5-enol	6.03%


**Fig S16:** Total ion chromatograms of GC-MS for the ethanol of *Orychophragmus violaceus* (L.) O. E. Schulz.

Table S16: GC-MS analysis result of ethanol extract of flowers from *Orychophragmus violaceus* (L.) O. E. Schulz.

Number	materials	relative content
1	Trifluoroethanol, TBDMS derivative	9.94%
2	Methanone, (4-bromo-5-methyl-2-nitro-3-thienyl)(phenyl)-	2.00%
3	Benzoic acid, 2-(dimethylamino)ethyl ester	1.59%
4	Sedoheptulosan	0.85%
5	Dodecane, 1-fluoro-	4.33%
6	3-n-Butylthiolane	0.64%
7	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	28.02%
8	3-n-Butylthiolane	11.23%
9	6-O-Methyl-2,4-methylene-.beta.-sedoheptitol	0.44%
10	Methyl 3,5-tetradecadiynoate	0.08%
11	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	1.41%
12	2,5-Octadecadiynoic acid, methyl ester	1.56%
13	3,6-Diazahomoadamantan-9-one Hydrazone	3.98%
14	2,6-Di-tert-butyl-4-hydroxy-4-methylcyclohexa-2,5-dien-1-one	1.82%
15	5,8,11-Eicosatrienoic acid, (Z)-, TMS derivative	0.27%
16	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	4.15%
17	10-Undecynoic acid, TMS derivative	4.63%
18	1,25-Dihydroxyvitamin D3, TMS derivative	1.05%
19	Estra-1,3,5(10)-trien-17.beta.-ol	7.09%
20	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	1.78%
21	Hexadecanoic acid, ethyl ester	13.15%




Fig S17: Total ion chromatograms of GC-MS for the ethanol of *Leucanthemum maximum* (Ramood) DC.

Table S17: GC-MS analysis result of ethanol extract of *Leucanthemum maximum* (Ramood) DC.

Number	materials	relative content
1	Trifluoroethanol, TBDMS derivative	6.11%
2	Benzamide, N-[6-(2-furyl)-2-oxo-2H-pyran-3-yl]-	1.27%
3	Benzoic acid, 2-(dimethylamino)ethyl ester	0.22%
4	Methcathinone	0.27%
5	(Methoxymethyl)trimethylsilane	2.01%
6	3-n-Butylthiolane	1.23%
7	Methyl.alpha.d-rhamnopyranoside	0.16%
8	(2-Benzyloxy-2-oxiran-2-ylethoxy)-t-butyldimethylsilane	0.24%
9	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	3.17%
10	2,5-Octadecadiynoic acid, methyl ester	0.89%
11	.alpha.,.beta.-Gluc-octonic acid lactone	0.51%
12	Methyl 4,7,10,13-hexadecatetraenoate	1.17%
13	cis-5,8,11,14,17-Eicosapentaenoic acid	0.89%
14	2,2-Dimethyl-5-[2-(2-trimethylsilylethoxymethoxy)-propyl]-[1,3]dioxolane-4-carboxaldehyde	0.65%
15	9,12-Octadecadiynoic acid, trimethylsilyl ester	0.89%
16	Linoleic acid ethyl ester	13.41%
17	9,12-Octadecadiynoic acid, tert-butyldimethylsilyl ester	0.83%
18	11-Dodecenoic acid, 10-hydroxy-, methyl ester	3.55%

Number	materials	relative content
19	10-Undecynoic acid, TMS derivative	17.27%
20	bis[(4E)-Hex-4-en-1-yloxy](dimethyl)silane	1.48%
21	Methyl 5,7-hexadecadiynoate	1.64%
22	1,25-Dihydroxyvitamin D3, TMS derivative	1.21%
23	9,12-Octadecadienoic acid (Z,Z)-	5.55%
24	5,8,11-Eicosatriynoic acid, methyl ester	7.20%
25	5,8,11,14-Eicosatetraynoic acid	17.63%
26	Estra-1,3,5(10)-trien-17.beta.-ol	3.49%
27	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	4.75%
28	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	1.40%
29	5,8,11,14-Eicosatetraynoic acid, TMS derivative	0.91%




Fig S18: Total ion chromatograms of GC-MS for the ethanol of *Oenothera speciosa* Nutt.

Table S18: GC-MS analysis result of ethanol extract from *Oenothera speciosa* Nutt.

Number	materials	relative content
1	10,12-Octadecadiynoic acid	0.73%
2	Perfluoropropionic acid, TBDMS derivative	4.72%
3	(1S,2S)-(+)-N-Methylpseudoephedrine	1.22%
4	Ephedrine	0.19%
5	Methoxymethyl)trimethylsilane	1.45%

Number	materials	relative content
6	Benzoyl isothiocyanate	0.33%
7	4-Cyclohexene-1,2-diol	0.30%
8	2,5,6-Trimethyl-1,3-oxathiane	1.12%
9	3-n-Butylthiolane	4.85%
10	5-Thio-D-glucose	2.56%
11	Methyl 6-oxoheptanoate	0.94%
12	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	4.91%
13	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	19.99%
14	5,8,11,14-Eicosatetraynoic acid	6.54%
15	t-Butyl(4,5-dihydrofuran-2-yl)dimethylsilane	0.47%
16	2,2-Dimethyl-5-[2-(2-trimethylsilylethoxymethoxy)-propyl]-[1,3]dioxolane-4-carboxaldehyde	0.22%
17	Melezitose	1.34%
18	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	1.39%
19	4-Hexenoic acid, 4-methyl-6-(fluorodimethylsilyl)-6-trimethylsilyl-	6.36%
20	Benzene, 5-heptene-1,3-diyn-1-yl-	36.79%
21	10-Undecynoic acid, TMS derivative	0.65%
22	1,25-Dihydroxyvitamin D3, TMS derivative	0.22%
23	9,10-Secosteroid-5,7,10(19)-triene-3,24,25-triol, (3.beta.,5Z,7E)-	0.82%
24	Phen-1,4-diol, 2,3-dimethyl-5-trifluoromethyl-	1.89%




Fig S19: Total ion chromatograms of GC-MS for the ethanol of *Iris tectorum* Maxim.

Table S19: GC-MS analysis result of ethanol extract from *Iris tectorum* Maxim.

Number	materials	relative content
1	Perfluoropropionic acid, TBDMS derivative	6.50%
2	Trifluoroethanol, TBDMS derivative	2.45%
3	Ephedrine	1.24%
4	1,3-Propanediol, TMS derivative	0.59%
5	Benzoyl isothiocyanate	0.33%
6	Tetraethyl silicate	0.81%
7	3-n-Butylthiolane	4.77%
8	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	9.09%
9	Methyl 6-oxoheptanoate	2.65%
10	tert-Nonyl mercaptan	0.15%
11	2,2-Dimethyl-5-[2-(2-trimethylsilylethoxymethoxy)-propyl]-[1,3]dioxolane-4-carboxaldehyde	1.87%
12	10-Undecynoic acid, TMS derivative	3.76%
13	Melezitose	10.41%
14	1-Tributylsilyloxy-3-phenylpropane	1.76%
15	Tetradecanoic acid	5.19%
16	Tetradecanoic acid, ethyl ester	11.82%
17	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	2.21%
18	tert-Hexadecanethiol	10.00%
19	Estra-1,3,5(10)-trien-17.beta.-ol	5.29%
20	Hexadecanoic acid, ethyl ester	18.46%
21	9,10-Secocholesta-5,7,10(19)-triene-3,24,25-triol, (3.beta.,5Z,7E)-	0.65%




Fig S20: Total ion chromatograms of GC-MS for the ethanol of *Gaillardia pulchella* Foug.

Table S20: GC-MS analysis result of ethanol extract from *Gaillardia pulchella* Foug.

Number	materials	relative content
1	10,12-Octadecadiynoic acid	10.49%
2	Perfluoropropionic acid, TBDMS derivative	2.71%
3	Methanone, (4-bromo-5-methyl-2-nitro-3-thienyl)(phenyl)-	0.69%
4	Benzoic acid, 2-(dimethylamino)ethyl ester	2.18%
5	Methoxymethyltrimethylsilane	5.80%
6	1,3-Propanediol, 2-methyl-, dipropionate	1.18%
7	Nonanoic acid	0.18%
8	2-Dimethylsilyloxytridecane	0.56%
9	(E)-3,7-Dimethyl-2,6-octadien-1-yl propionate	0.32%
10	3-Azabutyl-1-ol, 4-cyclopropyl-3,3-dimethyl-, bromide	0.39%
11	Dodecane, 1-fluoro-	0.71%
12	(+)-cis-Verbenol, trimethylacetate	0.26%
13	Eucalyptol	1.82%
14	3-trsns-(1,1-dimethylethyl)-4-trans-methoxycyclohexanol	0.13%
15	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	5.70%
16	3-n-Butylthiolane	1.01%
17	.alpha.-Linolenic acid, TMS derivative	1.94%
18	Pregnane-3,20-diol, (3.alpha.,5.beta.,20S)-, 2TMS derivative	2.82%
19	Melezitose	3.79%

Number	materials	relative content
20	.beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimidate]	0.31%
21	2,5-Octadecadiynoic acid, methyl ester	3.74%
22	(1R,2S,6S,7S,8S)-8-Isopropyl-1-methyl-3-methylenetricyclo[4.4.0.02,7]decane-rel-	1.79%
23	[1-(3,3-Dimethyloxiran-2-ylmethyl)-3,7-dimethylocta-2,6-dienyl]trimethylsilane	0.96%
24	[1,1'-Bicyclopropyl]-2-octanoic acid, 2'-hexyl-, methyl ester	4.34%
25	9,12-Octadecadiynoic acid, tert-butyldimethylsilyl ester	1.48%
26	Octadecanoic acid, ethyl ester	3.91%
27	2,2-Dimethyl-6-methylene-1-[3,5-dihydroxy-1-pentenyl]cyclohexan-1-perhydrol	7.54%
28	2,5-Octadecadiynoic acid, methyl ester	0.55%
29	Estra-1,3,5(10)-trien-17.beta.-ol	22.01%
30	7-Methyl-Z-tetradecen-1-ol acetate	7.92%
31	Arachidonic Acid, TBDMS derivative	0.24%
32	5,8,11-Eicosatrienoic acid, (Z)-, TMS derivative	2.30%
33	t-Butyl-(2-[3-(2,2-dimethyl-6-methylene-cyclohexyl)-propyl]-[1,3]dithian-2-yl)-dimethyl-silane	0.23%

Publisher's Note: Engineered Science Publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits the use, sharing, adaptation, distribution and reproduction in any medium or format, as long as appropriate credit to the original author(s) and the source is given by providing a link to the Creative Commons license and changes need to be indicated if there are any. The images or other third-party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.