

Multi-Analytical and In Silico Evaluation of Commercial Liquid Smokes: Volatile Profiling, Polycyclic Aromatic Hydrocarbons Safety Assessment and Molecular Mechanisms against Deoxyribonucleic Acid Gyrase B

Yusnaini,^{1,2,*} Sri Lestari,¹ Khusna Arif Rakhman,³ Muhammad Ikhlas Abdjan,⁴ Fadjar Mulya^{5,6} and Eka Kusuma Dewi⁷

¹Faculty of Agriculture, University of Khairun, Ternate, North Maluku, 97728, Indonesia

²Faculty of Agriculture, University of Khairun, Ternate, North Maluku, 97728, Indonesia

³Department of Chemistry Education, University of Khairun, Ternate, North Maluku, 97728, Indonesia

⁴Department of Chemistry, Faculty of Mathematics and Natural Sciences, University of Negeri Surabaya, Surabaya, East Java, 60213, Indonesia

⁵Department of Engineering, Faculty of Advanced Technology and Multidiscipline, University of Airlangga, Surabaya, East Java, 60115, Indonesia

⁶Computational Health and Medicine Research Group, Institute of Tropical Disease, University of Airlangga, Surabaya, East Java, 60115, Indonesia

⁷Faculty of Agriculture, University of Khairun, Ternate, North Maluku, 97728, Indonesia

*Email: yusnaini@unkhair.ac.id (Yusnaini)


Received: 04 May 2026; Revised: 25 June 2026; Accepted: 15 October 2026

Type: Research article.

Abstract

Liquid smoke is increasingly utilized as a natural food preservative; however, the chemical standardization and sub-atomic mechanisms of commercial brands remain inadequately understood. This study evaluates the physicochemical characteristics, volatile profiles, and antioxidant/antibacterial potencies of three commercial liquid smoke brands (LS1, LS2, and LS3) through an integrated experimental and computational framework. Gas Chromatography-Mass Spectrometry (GC-MS) analysis identified 59 volatile constituents, revealing distinct chemical fingerprints ranging from cellulose-derived levoglucosan in LS1 (49.69%) to phenolic-rich profiles in LS3 (phenol: 42.31%). Safety assessments highlighted that while all brands demonstrated efficacy, only LS1 contained detectable levels of the regulated carcinogen benzo[a]pyrene (6.35 µg/L), whereas LS3 exhibited the cleanest toxicological profile. Experimentally, LS1 showed the highest antioxidant potency ($IC_{50} = 38.44 \mu\text{g/L}$) and antibacterial efficacy against *E. coli*, *S. aureus*, and *B. cereus* (inhibition zones > 25 mm). Density Functional Theory (DFT) calculations corroborated these findings, showing that the superior reactivity of 2,4-dimethoxyphenol (C5) is driven by a narrow HOMO-LUMO energy gap (5.20 eV) and low global hardness ($\eta = 2.60$). Furthermore, molecular docking confirmed that these volatiles act as multi-target inhibitors, with 2-furanmethanediol dipropionate (C1) and C5 showing high binding affinities (Grid Scores: -40.49 and -36.06 kcal/mol) for the ATP-binding pocket of Deoxyribonucleic Acid Gyrase B.

Table of Contents



Innovative Description: This study evaluates the chemical properties, flavor profiles, and biological activities of three commercial liquid smoke brands using experimental and computational methods.